

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehponline@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Population-Level Exposure to Particulate Air Pollution during Active Travel: Planning for Low-Exposure, Health-Promoting Cities

Steve Hankey, Greg Lindsey, and Julian D. Marshall

Table of Contents

Summary of land use variables and output of spatial models

Table S1. Summary of land use variables

Table S2. Descriptive statistics of model outputs

“Sweet-spot” neighborhoods by pollutant

Figure S1. Neighborhood-types for particle number concentration. Maps were created in ArcMap; underlying street and land use data are from <https://gisdata.mn.gov/>.

Figure S2. Neighborhood-types for black carbon concentration. Maps were created in ArcMap; underlying street and land use data are from <https://gisdata.mn.gov/>.

Figure S3. Neighborhood-types for PM_{2.5} concentration. Maps were created in ArcMap; underlying street and land use data are from <https://gisdata.mn.gov/>.

Results of spatial models stratified by land use and transportation network variables

Figure S4. Bicycle and pedestrian traffic volumes and particulate concentrations stratified by road type. This plot includes off-street trails unlike Figure 3 in the main text.

Figure S5. Bicycle and pedestrian traffic volumes and particulate concentrations stratified by population density

Figure S6. Bicycle and pedestrian traffic volumes and particulate concentrations stratified by land use mix

Figure S7. Bicycle and pedestrian traffic volumes and particulate concentrations stratified by open space area

Figure S8. Bicycle and pedestrian traffic volumes and particulate concentrations stratified by retail area

Figure S9. Bicycle and pedestrian traffic volumes and particulate concentrations stratified by population density, land use mix, and open space area. Concentrations shown as absolute values instead of normalized values (Figure 4 in the main text)

Figure S10. Bicycle and pedestrian traffic volumes and particulate concentrations stratified by industrial area

Figure S11. Bicycle and pedestrian traffic volumes and particulate concentrations stratified by share of non-white residents

Figure S12. Bicycle and pedestrian traffic volumes and particulate concentrations stratified by household income

Comparison of afternoon rush-hour bicycle and pedestrian traffic with morning rush-hour particulate concentrations

Figure S13. Replication of Figure 1 (main text) using a morning rush-hour (7-9am) concentration surface. Spatial patterns of each factor (left-panel); plots of the transect (right-panel). The plots follow the transect from point A (left) to point B (right). Maps were created in ArcMap; underlying street and land use data are from <https://gisdata.mn.gov/>.

Figure S14. Replication of Figure 2 (main text) using a morning rush-hour (7-9am) concentration surface. The maps represent the four categories of neighborhood-type outlined in Table 2 of the main text. Maps were created in ArcMap; underlying street and land use data are from <https://gisdata.mn.gov/>.

Figure S15. Replication of Figure 3 (main text) using a morning rush-hour (7-9am) concentration surface. Trends reflect similar patterns to afternoon rush-hour with more blocks meeting the 15% reduction in BC or PN criteria for the morning surface. Maps were created in ArcMap; underlying street and land use data are from <https://gisdata.mn.gov/>.